

Help the Aged produces 24 free advice leaflets (see inside for full list). These are available from the Information Resources Team at the address below, email advicelaflets@helptheaged.org.uk or fax: 020 7239 1839.

January 2006

© Help the Aged CS/00 ID5233 01/06 Company limited by guarantee Registered in England no 1263446 Registered charity no 272786

Fighting for disadvantaged older people in the UK and overseas,

WE WILL:

COMBAT POVERTY wherever older people's lives are blighted by lack of money, and cut the number of preventable deaths from hunger, cold and disease

REDUCE ISOLATION so that older people no longer feel confined to their own home, forgotten or cut off from society

CHALLENGE NEGLECT to ensure that older people do not suffer inadequate health and social care, or the threat of abuse

DEFEAT AGEISM to ensure that older people are not ignored or denied the dignity and equality that are theirs by right

PREVENT FUTURE DEPRIVATION by improving prospects for employment, health and well-being so that dependence in later life is reduced

Head Office, 207–221 Pentonville Road, London N1 9UZ

T 020 7278 1114 F 020 7278 1116

E info@helptheaged.org.uk www.helptheaged.org.uk

HELPTHEAGED WE WILL™

Managing Your Medicines



Taking medication safely

Advice for older people

Endorsed by



Help the Aged produces a range of free advice leaflets for older people

Financial leaflets

- Can You Claim It?
- Check Your Tax
- Claiming Disability Benefits
- Questions on Pensions
- Thinking About Money

Housing and home safety leaflets

- Care Homes
- Fire
- Help in Your Home
- Housing Matters
- Keep Out the Cold
- Your Safety
- Your Security

Health leaflets

- Bereavement
- Better Hearing
- Better Sight
- Bladder and Bowel Weakness
- Fight the Flu
- Fitter Feet
- Healthy Bones
- Healthy Eating
- Keeping Mobile
- Managing Your Medicines
- Shingles
- Staying Steady

Leaflets are free of charge and available from the Information Resources Team at the address on the back page, email advicleaflets@helptheaged.org.uk or fax: 020 7239 1839.

‘Managing Your Medicines’ was written in association with Dr Peter Rivers of the Pharmacy Academic Practice Unit at the University of Derby. It is endorsed by the Primary and Community Care Pharmacy Network (PCCPN). This edition of the leaflet was updated by Dr Rivers and members of the PCCPN National Committee.

If you would like this leaflet in another format, such as large print or audio tape, please contact the Information Resources Team on **020 7278 1114**.

Contents

Getting medicines on prescription	page 4
Buying medicines	page 5
Medicine names	page 6
Getting the best from your medicines	page 6
Remembering to take your medicines	page 7
Safe-keeping	page 9
Expiry dates	page 9
Reducing side-effects	page 10
Falls	page 11
Other types of medicines	page 12
Reviewing your medicines	page 13
Disposing of medicines	page 14
Taking your medicines safely	page 15

We're all used to getting a prescription from the doctor when we're not well. It's a fact of life that, as we get older, we are more likely to need medicines prescribed by the doctor.

Most of the advice given in this leaflet applies to people of all ages. However, there are several reasons why older people should take extra care with medicines. For example, as we become older, we will probably need smaller doses of medicines. This is because our liver and kidneys become less efficient at removing medicines, so that some medicines stay active in the body for longer. Don't worry about this - it's quite normal. It just means that you must be very careful not to take more of a medicine than your doctor tells you to. This leaflet explains how to use your medicines safely and effectively.

Getting medicines on prescription

Once you reach 60 you don't have to pay for prescription medicines. Just fill in and sign the form on the back of the prescription that says that you are exempt from prescription charges. If you find filling the form difficult, ask a friend or relative to help you. You may be exempt from prescription charges, whatever your age, if you receive treatment for certain long-term conditions, such as diabetes. Ask your pharmacist for advice.

If you're under 60 and have to pay for your prescription, you might find it cheaper to buy a **pre-payment certificate**. This will cover all your prescription charges for varying periods of time. Application forms are available from most pharmacies.

When you receive a prescription, you should collect your medicines as soon as possible. Some pharmacists and doctors offer services to make collecting your prescriptions easier.

- Some pharmacies will collect your prescription from your doctor's surgery so it is ready to pick up from the pharmacy of your choice. Ask your pharmacist if they offer this service.
- If you are housebound and unable to get out it is worth asking if your pharmacist can deliver your prescription to you at home. Check whether or not there will be a charge for this.
- Some local voluntary agencies may also be able to collect your medicines for you.
- If your medicines have stayed the same for a long time you could use a repeat dispensing service. This means that you don't have to go to the surgery every time you need more medicines, you just have to go to the pharmacy. Ask your doctor for more information.

Buying medicines

It is important to check that any medicines and herbal medicines you buy are safe to take together and with any prescribed medicines you're taking. This is because different medicines and remedies may contain the same ingredients. For example, a medicine that your doctor has prescribed and one that you buy may both contain paracetamol. In this way, you could take more than the recommended daily dose of paracetamol without realising it.

You should also check that any medicines and herbal medicines you buy won't interfere with your prescribed medicines. Some medicines can interact with each other which could stop them from working properly or altogether. This could make you ill.

Always check with your pharmacist that any medicines and herbal medicines you buy are safe to take with your prescribed medicines.

Many pharmacists now keep records of medicines that are sold as well as dispensed to patients. This helps to make sure that the best possible advice is given.

Medicine names

Most medicines have two names. These are the **trade name** (or brand name) and the **approved name** (or ingredient name). The trade name is the one given to a medicine by its manufacturer. For example, imagine a fictional brand of coffee called 'Blue Valley'. Here, 'Blue Valley' is the trade name while 'coffee' is the approved name - the trade name is designed to be catchy and easy to remember.

With most medicines, it doesn't matter what particular brand (trade name) you use. With some medicines though, you need to keep to the same brand. This depends on a number of things - what your illness is, the range of medicines available, and your particular needs. Ask your doctor if you need to stick to the same brand. If you do, your doctor should prescribe using the brand name.

Getting the best from your medicines

- Always read the label on your medicines. If the instructions are not clear, ask your pharmacist or doctor to explain them. Ask your doctor to write the full instructions on your prescription, and not just 'as directed' as this can be confusing. The telephone number of your pharmacy may be on the label. You can ring this number to ask for advice about your medicines.
- The writing on medicine labels can be very small. Pharmacists may be able to provide large print labels and a few can also provide Braille labels for your medicines.

- Don't throw away the packaging from your medicines until you've finished taking them; and always read the patient information leaflet before starting to take your medicines.
- If the instructions on the label tell you to take the medicine on an empty stomach, this means that you should try to take the medicine at least an hour before, or two hours after, your meal.
- Always drink plenty of cool liquid with tablets and capsules - a full glass of water is ideal.
- Always take tablets and capsules when standing or sitting as upright as possible. This helps to stop them getting stuck in your throat.
- Don't crush tablets or open your capsules and dissolve them in a drink without checking with your pharmacist. This could stop them from working properly and in some cases can even be dangerous. If you find tablets difficult to swallow, ask your pharmacist or doctor if your medicine comes in a tablet that can be dissolved in water, or in a liquid form.
- Most tablet bottles have child-resistant caps, which can be very difficult to open, especially if you suffer from arthritis or can't get a strong grip on the container. You can ask your pharmacist for ordinary screw tops when handing in your prescription.

Remembering to take your medicines

Everyone finds it difficult to remember to take medicines. There are a few things you can do to make it easier. The main thing is to work out what times suit you best. Meal times are often a good time to take medicines because you're likely to have a drink with your meal. If you need to take your medicine on an empty stomach, try to get into a routine of taking the medicine at the same times each day.

Some people find it helpful to write down the times on a calendar and tick them off each time they take a dose. The pharmacist may be able to provide you with a chart to help remind you when to take your medicine.

Another way of remembering is to use a **memory aid container**. These are containers with small compartments that help you to organise your medicines so that you take them at the right time. There are many different types of memory aid container. For example, some containers have an alarm to remind you when to take your medicines. Others can have a recorded message from your doctor or pharmacist, telling you how many tablets to take and when to take them. Some open and close more easily than others, which can be important for someone with arthritis.

Ask your pharmacist for advice on the different types of memory aid container and how they work, or contact the **Disabled Living Foundation** on **0845 130 9177**.

You usually have to buy these memory aids yourself but you might find it's worth it if you're having problems remembering to take your medicines. Take your time choosing the best one for you. It is very important to make sure that you put the right medicines in the correct parts of the container. You could ask a friend, relative or carer to double-check against your prescription to be sure that the container is filled correctly. Take extra care if your prescription has changed recently. **If you are in any doubt, ask your pharmacist to check the contents again.**

Some people find it useful to draw up their own chart to help them remember to take their medicines. You could design your own headings. They could be things like:


- What time do I need to take my medicine?
- Do I need to take my medicine on an empty stomach?

You could have as many headings as you like, depending on the kind of things that you need to remind yourself of. Remember that you must update your chart if you ever change your medicines, and that you may need different instructions for different medicines. You could ask your pharmacist for advice.

Safe-keeping

Never share or give your medicines to anyone else. They have been prescribed for you and may be harmful to other people.

Never keep your medicines where children can reach them. Ideally medicines should be kept in a locked cupboard. If this isn't possible, make sure all your medicine containers are closed properly.



Expiry dates

Like foods, all medicines have a limited shelf-life called an expiry date or 'use-by' date. So it is best to get them in small quantities and to make sure you don't use them after the expiry date. There may be a date stamped on the container, or the letters EXP followed by a date. For example, EXP 1/12/07 would mean 'do not use after 1st December 2007'. If you can't find a date on the medicine container, as a general rule you shouldn't keep tablets or capsules for longer than a year. Liquids should only be kept for six months. If in doubt, check the expiry date with your pharmacist.

Some medicines go off very quickly once they're opened. The label will tell you not to use them after a specified amount of time. It is important to follow this advice.

Medicines will last longer if they are kept cool and dry. When choosing a place to store your medicines, try to avoid hot and steamy places like the bathroom or kitchen. Don't worry too much if the best place to store medicines is in the bathroom - but remember that steamy and warm conditions will reduce the shelf life.


The best way to avoid taking out-of-date medicines is to make sure you return them to your pharmacist once you've finished with them - don't hold on to them for a 'rainy day'.

Your pharmacist will make sure they are disposed of safely.

Reducing side-effects

There is no such thing as a medicine without any possible side-effects. Sometimes it might be worth putting up with some mild side-effects if it means that your infection or illness gets better. And, of course, some people may never experience any side-effects at all.

Stomach upsets are a common side-effect. You can reduce the risk of getting a stomach upset by taking your medicine while you're eating, or just after a meal. If you're not going to eat a main meal, take them with a



snack or milky drink. However, some medicines work better when taken on an empty stomach, say one hour before eating. If this is the case, the directions on the label will usually tell you to take before food.

If you think that you are experiencing side-effects you should mention it to your doctor even if it seems trivial. What you may feel is trivial could be important. It is easy to forget to ask the doctor about things like this, so it may help to write down the questions you want to ask before you go, to act as a reminder during your consultation.

If you do suffer from side-effects, the amount of medicine that you're taking might need adjusting, or your doctor might want you to try a slightly different medicine that is less likely to cause side-effects.

You should avoid alcoholic drink if you are taking certain types of medicine. Ask your pharmacist or doctor for advice on whether you can drink alcohol while on your medication.

If you do experience symptoms that you think might be side-effects from your medicines, get in touch with your doctor or pharmacist immediately.

Falls

Some medicines can make you feel drowsy or unsteady on your feet, and this might mean you're more likely to lose your balance and fall over. It's important to be aware that some medicines might make you feel like this. Take care when getting out of a chair or the bath. If you take sleeping tablets be extra careful if you have to get out of bed during the night and when you first get up in the morning. Feeling dizzy when you stand up is more common as you get older, and some medicines can make the problem worse.

Other types of medicines

You should take just as much care with other types of medicines, such as eye, ear and nose drops, inhalers and creams as with the medicines that you swallow.

Ear, eye and nose drops


Ear, eye and nose drops can cause side-effects just like tablets and capsules. Make sure you use the dose that is written on the container and **never share** your drops with anyone else. Once an eye drop container is opened, you shouldn't use it for longer than four weeks. This helps to prevent germs from getting into the eye drops.

You also need to be careful to stop infections being passed from one eye to the other. Try not to touch your eye when putting in your drops and if you have a separate container for each eye, make sure that you stick to using the same one each time. This will help prevent passing any infection from one eye to the other. If you want to know how to put in your eye drops, ask your pharmacist for advice.

If you find it difficult to put in your own drops, ask a friend, relative or carer for help, or think about using eye drop holders which can make things much easier. Your pharmacist can advise on these.

Creams and ointments

Creams and ointments should also be treated with care. **Always wash your hands** before and after applying a medicated cream (unless the cream is for your hands). If you are applying cream to someone else, protect your hands with disposable gloves. This is important. Although the cream is



meant to help someone else get better, it may contain ingredients that are harmful to you. Wearing gloves also helps prevent infections from passing between people.

When applying creams or ointments, you only need to use a small blob, and you should rub it in gently, unless the directions say otherwise.

Never offer your prescription cream or ointment or any other type of medicine to someone else - it could be harmful.

Inhalers

Inhalers are a very good way of getting medication straight into your lungs to help with your breathing. However, they can be difficult to use and some people don't manage to take in the right amount of medication.

If you have problems gripping and squeezing your inhaler, a special attachment can make the inhaler easier to use. Or you could use an inhaler which works as soon as you breathe in. Ask your pharmacist or doctor for more information.

If you have problems with your inhaler, or if it doesn't seem to be working properly, check with your pharmacist that you are using it correctly. If you still have problems, tell your doctor.

Reviewing your medicines

If you take medicines regularly, you might be able to get repeat prescriptions without seeing the doctor. **However, you must make sure that you still see your doctor regularly.** You may no longer need your medicine or the dose may need to be changed. Do tell your doctor if you think your medicines aren't working properly or could work better for you.

When you ask for a repeat prescription, only ask for the medicines you need. If you stock up on all your medicines at once, it's easy to get muddled if you have too many, or some might go out of date before you have a chance to use them.

If you think you've experienced any side-effects from your medicines, do mention it to your doctor or pharmacist, however unimportant you think it might be.

It's important to have your medicines checked every now and then by your doctor or pharmacist, to make sure that they're not out of date, or that none of them are interfering with your other medicines. You could take all your medicines with you when you go for your next doctor's appointment. You could also ask your pharmacist if they would go through your medicines with you.

Your medicines may need to be changed if your health changes or if you have to go into hospital. It is very important that you take all your medicines with you if you go into hospital, including any you have bought yourself. Having a complete list of medicines helps your doctor and pharmacist to make a proper assessment of your care.

Regular checks are essential so that you and your doctor can be sure that your medicines are working to your benefit. It's also important to check that you're not taking any medicines that are out of date or that interfere with any of your other medicines.

Disposing of medicines

If your medicines are out of date, or if you no longer need to take or use them, take them to a pharmacy for safe disposal.

Don't throw medicines out with household rubbish.

If you find it difficult to get out and about, ask a friend or relative to return the medicines to a pharmacy for disposal.

Taking your medicines safely

Do follow the instructions on the label. If you don't understand them, ask your pharmacist or doctor to explain.

Do take medicines at meal times to help you remember them (unless the directions tell you to take them on an empty stomach).

Do ask your pharmacist for advice. If you buy any medicines over the counter check that the ingredients do not duplicate or interfere with what you are already taking.

Do check whether you need to stick to the same brand of medicine when offered repeat prescriptions.

Do make regular appointments with your doctor to make sure that the medicines that have been prescribed for you are still appropriate.

Do make sure that you fully understand the directions that the doctor or pharmacist has given you.

Don't crush your tablets or open capsules and dissolve them in a drink without checking with your pharmacist or doctor first.

Never take more than the dose stated on the label of your medicine container.

Never share your medicines or give them to anyone else.

Avoid taking extra medicines in addition to those prescribed by the doctor.

Always take medicines you no longer need or use back to a pharmacy.